arXiv:gr-qc/0409007v3 25 May 2005

Counter example to a quantum inequality

S. Krasnikov

Abstract

A ‘quantum inequality’ (a conjectured relation between the energy density of a free quantum field and the time during which this density is observed) has recently been used to rule out some of the macroscopic wormholes and warp drives. I discuss the possibility of generalizing that result to other similar spacetimes and first show that the problem amounts to verification of a slightly different inequality. This new inequality *can* replace the original one, if an additional assumption (concerning homogeneity of the ‘exotic matter’ distribution) is made, and *must* replace it if the assumption is relaxed. Then by an explicit example I show that the ‘new’ inequality breaks down even in a simplest case (a free field in a simply connected two dimensional space). Which suggests that there is no grounds today to consider such spacetimes ‘unphysical’.

1 Introduction

Suppose a freely falling observer measures the renormalized expectation value of the energy density ρ of a free quantum field in a state ω. Denote

$$\overline{\rho}(\omega) = \mathcal{T}^{-1}\int_{\tau_1}^{\tau_2} \rho(\omega;\tau)\,\mathrm{d}\tau, \qquad \mathcal{T} \equiv |\tau_2 - \tau_1|$$

where the integral is taken along the observer's world line γ parametrized by the proper time τ. Obviously $\overline{\rho}$ is the average energy density. The quantity we are interested in is $\overline{\rho}\mathcal{T}^d$, where d is the dimensionality of spacetime. More specifically, we shall analyze the inequality

$$-\overline{\rho}\mathcal{T}^d \lesssim 1 \tag{1a}$$

(it is not trivial because we are concerned, in particular, with negative ρ) under the condition

$$\mathcal{T} \lesssim \mathcal{T}_R \equiv \left(\max|R_{\hat{\imath}\hat{\jmath}\hat{m}\hat{n}}|\right)^{-1/2}. \tag{1b}$$

In this expression the maximum is taken over all $\tau \in (\tau_1, \tau_2)$ and all sets of indices. The hats over the indices mean that the components of the Riemann tensor are found in the (orthonormal) frame associated with the observer.

In the next section I discuss the importance of inequality (1) in determining whether a spacetime is physically realizable. In section 3 the massless scalar field is considered in the two-dimensional de Sitter space. By explicit calculations the existence is demonstrated of a 'reasonable' state ω such that (1) is violated (in this letter I regard a state ω as reasonable if $\langle\omega|T_{ab}|\omega\rangle^{\mathrm{ren}}$ is smooth over the whole spacetime).

2 The quantum inequalities

Within the last decade dozens of papers (see [1] or [2] for a review) have been published on inequalities that look similar to (1) and express the idea that "the more negative the energy density is in an interval, the shorter must be the duration of the interval" [1, 3]. Unfortunately, all of them are often called equally — quantum inequalities (QI), which leads sometimes to confusion and even mistakes. So, let us first establish terminology.

Pick a non-negative function f normalized by

$$\int_{-\infty}^{-\infty} f(\tau)\,\mathrm{d}\tau = 1$$

and consider the 'weighted average' of the energy density

$$\rho_f(\tau_0;\omega) \equiv \int_{-\infty}^{-\infty} \rho(\tau;\omega) f(\tau-\tau_0)\,\mathrm{d}\tau$$

(both integrals are taken along γ, see the Introduction). Denote by $\mathcal{F}_{\omega_0}$ the Fock space built on a quantum state ω_0. Then *difference* QIs are the statements of the form

$$\rho_f(\omega) - \rho_f(\omega_0) \geq Q \qquad \forall\,\omega \in \mathcal{F}_{\omega_0}, \tag{I}$$

where Q does not depend on ω and τ_0. Their more general variety (note the last quantifier) is

$$\rho_f(\omega) - \rho_f(\omega_0) \geq Q \qquad \forall\,\omega \in \mathcal{F}_{\omega_0},\ \forall\,\omega_0. \tag{II}$$

A different type of QIs are those of the form

$$\rho_f(\omega) \geq Q \qquad \forall\,\text{reasonable}\ \omega. \tag{III}$$

Following [4] I shall call them *absolute.*

Almost all quantum inequalities established so far are of type I. Among them are, for example, those proved in [5] or in [6] (in the former ω_0 is the standard Minkowski vacuum ω_M and in the latter it is the vacuum defined by the timelike Killing field present in the spacetime). The second group is represented by the relation (4.1) of [7]. The absolute QI have been derived so far only for massless fields in two dimensions [8]–[10], as is stated in [4].

The distinction between the difference and the absolute QIs is *fundamental.* In particular, (I) does not become (III) even when $\rho_f(\omega_0) = 0$, which happens in the Minkowski space when $\omega_0 = \omega_M$, and perhaps may also occur in curved spacetimes for some special choice of ω_0. The reason is that the corresponding inequality holds not for the set of *all* (reasonable) ω, but only for a *very special subset* thereof (recall that ω can always be transformed into $\omega' \notin \mathcal{F}_{\omega_0}$ by an appropriate Bogolubov transformation). Thus a phrase like "…arbitrarily extended distributions of arbitrarily negative energy are not possible in Minkowski spacetime" [3] to be a *correct* inference from such a QI (the inequality [3, (1)] in this case) has to be complemented by quite a restrictive stipulation: "if the field is in the state $\omega \in \mathcal{F}_{\omega_M}$."

Remark. The procedure proposed in [4] (the transition to the limit $\Delta \to 0$, where Δ is the length of $\operatorname{supp} f$) transforms the inequality from (I) into one looking *similar* to that from (III) but not satisfying an essential condition $Q \neq Q(\omega)$. While the *real* (III) does not follow from (I) in this limit either.

The importance of the QIs lies basically in the fact that they supposedly "place some severe constraints on wormhole and warp drive spacetime geometries" [1]. Specifically, it was argued that the 'total negative energy' needed to maintain a human-size Alcubierre bubble or Krasnikov tube is about $10^{67}\,$g [11, 12] and that the 'absurdly benign' wormhole can exist only if the exotic matter supporting it is concentrated in a $10^{-68}\,$cm thick layer [13]. These estimates are obtained in assumption that for an appropriately chosen f with $\operatorname{supp} f \in (\tau_1, \tau_2)$

$$\rho_f(\omega) > -\mathcal{T}^{-d} \qquad \forall\, \mathcal{T} \lesssim \mathcal{T}_R, \quad \forall\, \text{reasonable } \omega. \tag{2}$$

in any spacetime for any (free) quantum field[1]. It should be emphasized that (2) is *a conjecture.* Being an absolute QI it, contrary to a common misconception, *neither follows from, nor is supported by any of multiple*

---

[1] In the original papers the assumption looked slightly different (in particular, f had infinite 'tails'). This form it took a little later [14]. There is also an additional requirement on $\mathcal{T}$, but in simply connected spacetimes without 'boundaries' it is always satisfied.

difference QIs, as discussed above. At present it can be substantiated *only* by Flanagan and Vollick's QIs for massless fields in two dimensions [8]–[10].

Once the assumption (2) is made the reasoning goes, roughly, as follows. Consider a point p through which there is a timelike geodesic segment $\gamma(\tau)$ of the length $\mathcal{T}_R$ with ρ negative and approximately constant along γ. Suppose that

$$\max|R_{\hat{\imath}\hat{\jmath}\hat{m}\hat{n}}(p)| \approx \max|T_{\hat{k}\hat{l}}(p)| \approx -\rho(p) \tag{3}$$

(p satisfying all these requirements happens to exist in all three above-mentioned spacetimes). Applying (2) with $\mathcal{T} = \mathcal{T}_R$ to γ one can write

$$|\rho_f| < \mathcal{T}_R^{-4} = \big(\max|R_{\hat{\imath}\hat{\jmath}\hat{m}\hat{n}}|\big)^2 \approx \rho^2(p) \tag{4}$$

or

$$|\rho(p)| \gtrsim \rho_f/\rho(p) \approx 1. \tag{5}$$

Thus the energy density in p must be Planckian! All the prohibitive values for the 'total amount of negative energy' that are interpreted as severe restrictions on the warp drives stem just from this fact.

Those restrictions *by themselves* are, however, of little interest: both warp drives were designed to illustrate the idea of 'faster-than-light' travel and to do that by as simple means as possible. It seems not improbable that it is just their simplicity that leads to the undesirable properties of the matter sources. That is why it would be important to find out to what spacetimes the foregoing reasoning can be generalized. A number of cases in which it does *not* work are already known [15, 13]. Among them are

i. The spacetimes in which the 'exotic matter' is generated by *interacting* quantum fields, because (2) *is known* to break down in this case, which is demonstrated with the Casimir effect;

ii. Some wormholes, because the condition mentioned in the footnote 1 does not hold there;

iii. The spacetimes in the relevant regions of which $\max|R_{\hat{\imath}\hat{\jmath}\hat{m}\hat{n}}(p)| \gg \max|T_{\hat{k}\hat{l}}(p)|$ for all observers (as, for example, in the vicinity of *any* gravitating body).

And yet, a lot of interesting spacetimes are conceivable that should be dismissed as 'unphysical' be (an analog of) the restriction (2) true.

Thus the quantum inequality (2) deserves to be tested most seriously. In doing so, however, one has to specify f more definitely, because generally $\rho_{f_1}(\omega) \neq \rho_{f_2}(\omega)$ (unless, of course, exotic matter is distributed homogeneously as in the example above, but no reasons are seen to restrict ourselves to this case). And it is quite possible that among QIs obtained from (2) by specifying f some are true and some other are not. For example, one can choose f to be the characteristic function of the interval (τ_1, τ_2):

$$f = h\colon \qquad h(\tau) = 1 \quad \text{for } \tau \in (\tau_1, \tau_2), \qquad h(\tau) = 0 \quad \text{otherwise.}$$

ρ_f then is just $\rho_h = \overline{\rho}$ and the inequality in (2) transforms into (1). As well one could choose another f:

$$f \in C^\infty, \qquad \int_\gamma f'^2 f^{-1}\, \mathrm{d}\tau < 1. \tag{6}$$

and get another inequality[2]. The state considered in the next section satisfies (2), (6) (in fully agreement with the results of [10]) and at the same time, as we shall see, violates (1).

Fortunately, one does not need to test *all* possible f. The point is that the whole reasoning cited above fails when so does (1). To see this consider a situation in which (1) breaks down, while (2) with $f = g \neq h$ does not. That clearly would imply

$$|\rho_g| \ll |\overline{\rho}|. \tag{7}$$

The approximate equality in (4) to hold it is necessary that $|\rho(p)| \approx \max_\gamma |\rho|$ and, as a consequence, $|\overline{\rho}| \lesssim |\rho(p)|$. The latter inequality combined with (7) implies $|\rho_g| \ll |\rho(p)|$. So, instead of (5) one gets

$$|\rho(p)| \gtrsim \rho_g/\rho(p) \ll 1.$$

Quite a useless result!

To summarize, if the inequality (1) held for all reasonable quantum states it would mean that a wide class of 'shortcuts' [13] require Planck scale energy densities. (1) in this implication can be substituted *neither* by a difference QI *nor* by the system (2),(6)[3].

---

[2] The similarity of these two inequalities does not, of course, make them the *same*. It is misunderstanding of this fact that underlies the recent paper [4]. For example in the key statement of the paper "we have shown that Krasnikov's supposed counterexample to the QEIs is, in fact, entirely compatible with them" the word 'QEIs' refers (or, at least, *must* refer) to inequality (1), while the word 'them' — to the inequality from (2) with f satisfying (6).

[3] Though if one day the latter is proven it will rule out the spacetimes in which exotic matter of moderate density is distributed homogeneously.

3 The counter example

That (1a) can be violated even in the Minkowski space follows immediately from the results of [8]. However, the corresponding state would have singularities at $\tau_{1,2}$. To construct a 'reasonable' state violating (1) consider the two-dimensional de Sitter spacetime. One of the customary choices of coordinates casts the metric into the form

$$\mathrm{d}s^2 = \frac{\alpha^2}{\sin^2(\mu-\nu)}\,\mathrm{d}\mu\mathrm{d}\nu, \qquad (\mu+\nu)\in \mathrm{I\!R},\ (\mu-\nu)\in(0,\pi).$$

However, to fix in a convenient way the quantum state of the field we shall use different coordinates. Pick a function w such that

$$w \in C^\infty(\mathrm{I\!R}), \quad w' \neq 0, \quad w(x)\big|_{(\epsilon,\frac{\pi}{2}-\epsilon)} = \tfrac{1}{2}\ln\tan x,$$

where ϵ is a small positive constant. [For example, w can be defined as

$$w(x) = \int_{\frac{\pi}{4}}^{x} [\chi_1(s) + \chi_2(s)\sin^{-1}2s]\,\mathrm{d}s,$$

where $\{\chi_i\}$ is a partition of unity for the cover $\{X_i\}$ of $\mathrm{I\!R}$, with $X_1 = (-\infty,\epsilon)\cup(\frac{\pi}{2}-\epsilon,\infty)$, and $X_2 = (\frac{\epsilon}{2},\frac{\pi-\epsilon}{2})$.] The new coordinates are defined as follows

$$u(\mu) = w(\mu), \quad v(\nu) = w(\nu), \qquad t = \tfrac{1}{2}(v-u), \quad x = \tfrac{1}{2}(v+u)$$

and the metric takes the form

$$\mathrm{d}s^2 = C\mathrm{d}u\mathrm{d}v, \tag{8a}$$

where C is smooth and nonzero in the whole space. In the region W, see figure 1,

$$W\colon \qquad \epsilon < \mu,\nu < \tfrac{\pi}{2}-\epsilon, \quad \nu<\mu,$$

C can be expressed explicitly:

$$C = \frac{\alpha^2}{\sin^2(\mu-\nu)}\frac{\mathrm{d}\mu}{\mathrm{d}u}\frac{\mathrm{d}\nu}{\mathrm{d}v}\Big|_W = \frac{4\alpha^2\cos^2\nu\cos^2\mu}{\sin^2(\mu-\nu)\cdot e^{-2u}e^{-2v}} = \alpha^2\sinh^{-2}(u-v). \tag{8b}$$



Figure 1: The region W (dark gray) of the de Sitter space (light gray) in different coordinates.

Consider now the massless scalar field in the conformal vacuum state associated with the metric (8). Its renormalized stress-energy tensor in W is readily found by formula (6.136) of [16]:

$$\begin{aligned} T_{vv} = T_{uu} &= -\tfrac{1}{12\pi}C^{1/2}\partial_u^2 C^{-1/2} = -\tfrac{1}{12\pi}. \\ T_{uv} &= \tfrac{1}{96\pi}RC = \tfrac{1}{12\pi}\sinh^{-2}(u-v) \end{aligned}$$

Let $\gamma \subset W$ be a geodesic segment

$$x = 0, \quad t \in [t_1, t_2],$$

where $t_{1,2}$ are chosen so that

$$\tfrac{1}{2}\ln\tan\epsilon < t_1 < t_2 \ll -1. \tag{9}$$

For an observer whose world line is γ the energy density is[4]

$$\rho = T_{\hat{0}\hat{0}} = T_{tt}C^{-1} = 2(T_{uu}-T_{uv})C^{-1} = -\tfrac{1}{6\pi}(1+\sinh^{-2}2t)C^{-1} \approx -(6\pi C)^{-1},$$

which yields

$$-\overline{\rho} = \frac{\mathcal{T}^{-1}}{6\pi}\int_{t_1}^{t_2} C^{-1}C^{1/2}\,\mathrm{d}t \approx \frac{\mathcal{T}^{-1}}{24\pi}\alpha^{-1}e^{-2t_1}.$$

At the same time

$$\mathcal{T} = -\int_{t_1}^{t_2}\alpha\sinh^{-1}2t\,\mathrm{d}t \approx \alpha\int_{2t_1}^{2t_2} e^s\mathrm{d}s \approx \alpha e^{2t_2}.$$

---

[4] I am grateful to Dr. Fewster for the detailed evaluation [4] of the coefficient (it is 1 and not $-1/2$ as was erroneously written in the draft of this letter) with which the term $\sinh^{-2}2t$ enters the expression. This coefficient, though, plays absolutely no role in what follows as we neglect the whole term [on the strength of the condition (9)].

Thus,

$$-\overline{\rho}\mathcal{T}^2 = \tfrac{1}{24\pi}e^{2(t_2-t_1)},$$

which obviously — just by picking sufficiently small ϵ and t_1 — can be made arbitrarily large in contradiction with (1) [note that

$$\mathcal{T} \ll \mathcal{T}_R = \alpha$$

and so the criterion (1b) is satisfied].

4 Conclusions

In a number of papers warp drives and wormholes have been labelled as unphysical because of the enormous energy densities supposedly needed for their existence[5]. This idea is supported by the results obtained for the Alcubierre bubble and Krasnikov tube and based on the conjecture (2),(6). The generalization of these results, however, would require the validity of a slightly different inequality, namely (1), for all reasonable quantum states in reasonable spacetimes. In fact, however, the inequality turns out to be invalid even in a simplest case of the massless scalar field in the two dimensional de Sitter space. Consequently, there are no reasons any longer to expect that the shortcuts are necessarily connected with Planckian energy densities. So, these objects are perhaps not that unphysical after all.

Acknowledgements

I wish to thank the members of the Friemann seminar and particularly Yu. Pavlov for helpful discussion.

References

[1] T. A. Roman, *gr-qc/0409090.*

[2] M. J. Pfenning, *gr-qc/9805037.*

---

[5] Ford and Roman argue that a large discrepancy in the length scales characterizing these spacetimes (if it takes place) would also be grounds for rejecting them as unphysical. But in the absence of explanation this thesis does not look convincing. The stars, for example, are nothing but associations of protons (and electrons), which are characterized by the length scale $r_{\rm pr} \sim 10^{-15}\,$m. So, Deneb, say, with its radius $R_{\rm D} \gtrsim 10^{11}\,$m is characterized by the length scales with the ratio $R_{\rm D}/r_{\rm pr} \gtrsim 10^{26}$. The ratio of the mass scales is $M_{\rm D}/m_{\rm pr} \gtrsim 10^{58}$. Is Deneb unphysical?

[3] L. H. Ford and T. A. Roman *Phys. Rev. D* **53** (1996) 5496.

[4] C. J. Fewster, *gr-qc/0409043*.

[5] L. H. Ford and T. A. Roman *Phys. Rev. D* **51** (1995) 4277.

[6] M. J. Pfenning and L. H. Ford *Phys. Rev. D* **57** (1998) 3489.

[7] C. J. Fewster *Class. Quantum Grav.* **17** (2000) 1897.

[8] É. É. Flanagan *Phys. Rev. D* **56** (1997) 4922.

[9] D. N. Vollick *Phys. Rev. D* **61** (2000) 084022.

[10] É. É. Flanagan *Phys. Rev. D* **66** (2002) 104007.

[11] M. J. Pfenning and L. H. Ford *Class. Quantum Grav.* **14** (1997) 1743.

[12] A. E. Everett and T. A. Roman *Phys. Rev. D* **56** (1997) 2100.

[13] S. Krasnikov *Phys. Rev. D* **67** (2003) 104013.

[14] L. H. Ford, M. J. Pfenning, and T. A. Roman *Phys. Rev. D* **57** (1998) 4839.

[15] K. D. Olum and N. Graham *Phys. Lett. B* **554** (2003) 175.

[16] N. D. Birrel and P. C. V. Davies, *Quantum Fields in Curved Space* (Cambridge: Cambridge University Press, 1982).